


5 YEAR RETROSPECTIVE STYLAGE® SATISFACTION SURVEY


Objective and method

- Customer satisfaction survey conducted between October 2012 and May 2013
- 80 practitioners, 8 countries, representing a population of more than 14,000 patients
- A retrospective questionnaire was sent to the practitioners using Stylage[®], asking for information about:
 - Patient population
 - o Aesthetic medical and technical practice
 - Product performance

- $\circ \, Comparison \, \, vs \, \, competitor \, products \, \,$
- o Immediate and delayed tolerance


Population






- About 50% of respondents declared having > 2 year follow-up of more than 100 patients
- Average years of experience of these practitioners with STYLAGE® products: 3 years (max 5 years, min 0.83 year)
- Representing > 8,000 patients followed-up for more than 2 years and > 6,000 from 1 to 2 years

Performance and satisfaction








Satisfaction Vs competitors according to 4 parameters

1 to 10 scale where 5 means = to competitors; Mean and CI95%

- 91% of respondents declared that Stylage® allows obtaining a better natural result than competitor products (score ≥6/10); 87% declared that it has both better volumizing and better lifting/smoothing capacities; 85% that it allows obtaining a better skin quality.
- Indeed, 70%, 56%, 61%, and 56% gave a score ≥8/10 in favour of Stylage range Vs competitors concerning natural result, volumizing capacity, lifting/smoothing effect, and skin quality respectively.

Practices

Do you use cannula in your daily practice?



Do you use products with lidocaine and/or perform anaesthesia in your daily practice?

| Product with lidocaïne and perform anaesthesia | 15% |
|--|-----|
| Product with lidocaïne alone | 21% |
| Anaesthesia alone | 36% |
| Neither of them | 9% |
| NA | 19% |

36% of the respondents used Stylage® products with lidocaine in their daily practice

What Stylage® products do you use in your daily practice?

Hydration (Hydro, Hydromax):

| | Other | Surgeon | Dermatologist | Cosmetic Doctor | Total |
|-------|-------|---------|---------------|-----------------|-------|
| No | 13% | 50% | 19% | 24% | 28% |
| Yes | 88% | 40% | 81% | 76% | 70% |
| NA | 0% | 10% | 0% | 0% | 3% |
| Total | 100% | 100% | 100% | 100% | 100% |

Fillers (Lips, S, M, L):


| | Other | Surgeon | Dermatologist | Cosmetic Doctor | Total |
|-------|-------|---------|---------------|-----------------|-------|
| No | 25% | 10% | 0% | 0% | 5% |
| Yes | 75% | 85% | 100% | 100% | 94% |
| NA | 0% | 5% | 0% | 0% | 1% |
| Total | 100% | 100% | 100% | 100% | 100% |

Volumizers (XL, XXL):

| | Other | Surgeon | Dermatologist | Cosmetic Doctor | Total |
|-------|-------|---------|---------------|-----------------|-------|
| No | 13% | 20% | 11% | 28% | 19% |
| Yes | 88% | 75% | 89% | 72% | 80% |
| NA | 0% | 5% | 0% | 0% | 1% |
| Total | 100% | 100% | 100% | 100% | 100% |

The most used Stylage[®] products are, in order, M, L, XL

Tolerance/Safety



Immediate Tolerance (in the 2 first weeks):

Average score: 8.9/10

• 91% of respondents gave ≥8/10

40.5% gave 10/10

Delayed Side Effects (more than 2 weeks after injection):

- 84% of respondents have never experienced any
- Global incidence of 0.025%, i.e.:
 - o 0.08 per 1,000 patients and per year
 - o 0.022 case per practitioner and per year

Limitations

- No individual patient data ⇒ but global practitioners' experience feedbacks
- Voluntary responding and no statistical testing
- The number of respondents by country is not representative of each market

